



CNRM, UMR 3589

SEMINAIRE CNRM N° 2020_07 mercredi 24 juin 2020 à 10h

Memory properties in cloud-resolving simulations of the diurnal cycle of deep convection par Chimène DALEU (METOFFICE)

en visioconférence

Visio conférences par Blue jeans : https://bluejeans.com/740523609/1839

Phone Dial-in +33.1.84.88.79.52 (France (Paris, French)) ID de la Réunion: 740 523 609 Code d'accès participants: 1839

<u>Abstract</u> :

A series of high-resolution three-dimensional simulations of the diurnal cycle of deep convection over land are performed using the new Met Office NERC cloud-resolving model. A memory function is defined to identify the effects of previous convection in modifying current convection. It is based on the probability of finding rain at time t0 given rain at previous time $t0 - \Delta t$ compared to the expected probability given no memory. The memory function is examined as a function of the lag time Δt . It is strongest at grey-zone scales (e.g., 5–10 kilometre), there is a change of behaviour for spatial scales between 10–15 kilometre, and it is reduced substantially for spatial scales larger than 25 kilometre. At grey-zone scales, there is a first phase of the memory function which represents the persistence of convection and it is maintained for about an hour. There is a second phase which represents the suppression of convection in regions which were raining 1 to 3 hours previously, and subsequently a third phases of the memory function develop earlier for weaker forcing.

When thermodynamic fluctuations resulting from the previous diurnal cycle are allowed to influence the development of convection on the next day, there are fewer rainfall events with relatively large sizes, which are more intense and thus, decay and recover more slowly, in comparison to the simulations where feedback from previous days is removed. Further sensitivity experiments reveal that convective memory attributed to these thermodynamic fluctuations resides in the lower troposphere.