

Radiation Experiments and Developments



Emily Gleeson¹, Laura Rontu², Kristian Pagh Nielsen³, Velle Toll⁴
¹Met Éireann, Ireland, ²FMI, Finland, ³DMI, Denmark, ⁴Reading University, UK

contact e-mails: emily.gleeson@met.ie, laura.rontu@fmi.fi, kpn@dmi.dk, v.toll@reading.ac.uk

Summary

What's new since the ASM in Lisbon?

- MarcoPolo aerosol experiments (FP7) [1]
- CIRC comparisons of radiation schemes
- Comparison of Tegen vs CAMS aerosols
- Working version of MUSC cycle 43
- Calling radiation subroutines intermittently vs calling them every time step
- Validation of HLRADIA using FMI archived operational data [2]

1. MarcoPolo Experiments

- As part of the FP7 project [1]: “MarcoPolo” aerosol experiments were run for a domain over China around Shanghai
- **Experiment 1:** HARMONIE-AROME cy40h1 default version
- **Experiment 2:** As above + MACC reanalysis aerosols converted to IFS aerosol categories
- **Experiment 3:** As above + Menon et al.. aerosol CCN/ τ_e , liq. indirect effect
- **Strong impacts on convective events seen in experiments 2 & 3.** This is mainly due to the strong increase in urban aerosols which affects the temperatures (Fig. 1 & 2)




Figure 1: Changes (Exp. 2 - Exp. 1) in liquid precipitation due to the direct aerosol effect of MACC reanalysis aerosols.





Figure 2: Changes (Exp. 3 - Exp. 2) in precipitation due to the indirect CCN effect of MACC reanalysis aerosols.

2. CIRC Experiments

- CIRC: Continual Intercomparison of Radiation Codes
- <http://circ.gsfc.nasa.gov/>
- Considered the 9 CIRC test cases show in the table
- Compared output from the HLRADIA, NBM (narrow band model) and LBL (line by line model, CHARTS) [2]

CIRC Experiment	
1b:	dry atmosphere, clear
2b:	very humid atmosphere, clear
3b:	humid atmosphere, clear
4b:	albedo=0.67, very dry atmosphere, clear
5b:	Same as 4b but with 2 x CO ₂
6b:	thick overcast liquid cloud, humid atmosphere
6d:	As 6b but clear sky.
7a:	moderately thin overcast liquid cloud, humid atmosphere
7b:	As 7a but clear sky


- **Clear-sky SW flux** is overestimated by HLRADIA at the surface and TOA (6-19 W/m²) and atmospheric absorption is underestimated
- **Cloudy-sky SW flux** at TOA overestimated by HLRADIA (~20 W/m²) – sensitive to how cloud droplet size is treated



- **Clear-sky LW flux** errors are small at the surface (within 7 W/m²). **Cloudy-sky LW flux** errors are small and positive (tuning needs investigation)
- HLRADIA strongly **overestimates the cloud LW radiative effect** at TOA
- Clouds with separate cloud layers – HLRADIA ok for SW but problems in the LW as the scheme accounts for clouds as a single layer but in reality there are strong exchanges between cold high clouds and warm low clouds

3. Radiation verification: CSI

- Using measured SW fluxes to verify modelled clouds is an improved method of verification compared to using synoptic surface observations
- In the latter only cloud cover is verified, whereas downwelling SW fluxes are an indirect measure of cloud water load and cloud microphysical properties
- We used the clear sky index (CSI) as a metric for SW flux and cloud verification (CSI is the global SWD radiation normalised by the estimated clear sky downwelling SW radiation) [3]
- Observations from 7 stations in Ireland were used in the verification (Fig. 4)



- Using CSI as a proxy for cloudiness highlights the binary (on/off) cloud cover in HARMONIE-AROME (Fig. 4)
- From a radiation view-point the differences between cycle 38 and 40 include:
 - Inhomogeneity factor (0.7 vs 1.0)
 - Nielsen cloud liquid optical properties
 - HARATU

4. CAMS Aerosols

- There are now 2 aerosol climatologies available in HARMONIE-AROME: Tegen (default [4]) and CAMS (Copernicus Atmosphere Monitoring Service)
- CAMS: AOD at 550 nm was derived using data from 2003-2011
- Relative to Tegen, CAMS land aerosols have a lower AOD over Northern Europe; the sea aerosols have a higher AOD (Fig. 5)






Fig. 6 shows the difference in global radiation and integrated cloud water when Tegen is replaced by CAMS aerosols



5. Frequency of call to radiation physics routines

- In mesoscale models fast interactions between clouds and radiation and the surface and radiation can be of greater importance than accounting for the spectral details of clear-sky radiation



- Fig 7: an example of the influence of the frequency of calling the IFS radiation scheme in a HARMONIE-AROME experiment
- Differences in average SW (left) and LW (right) downward surface fluxes over 1 hour from 0 to 1 UTC (8-9 am local time) on the 30th of July 2010 are shown
- Flux differences: radiation call every 15th time-step (default) minus radiation call every time-step